



IN THE CLAIMS

1. (Currently Amended) A steroid derivative selected from the group of compounds defined by formula (I) or (II) as shown below, wherein the only difference between said formulas is the bond between carbon number 5 and carbon number 6:



(I)



(II)

wherein

R_1O is in the β -position and R_1 is a hydrogen atom; ~~an NO_2 , an SO_3H , an $OP(OH)_3$ an acyl group, or any other group that forms an ester with an inorganic or organic acid; a protecting group, in the form of~~ such as CH_3 , CH_2OMe , or CH_2O -alkyl; ~~an aliphatic chain which is straight or branched, saturated or unsaturated, or cyclic, including mixed cyclic and aliphatic substituents, which substituents are saturated or unsaturated, aromatic or heterocyclic and contains up to 20 carbon atoms, which substituents can be chosen from hydroxyl, any halogen, amino or alkylamino, carboxylic acid or carboxylic acid ester;~~

R_2 is $R'O$ in β -position of carbon number 7 or can be $[[is]]$ hydrogen ~~in the case of formula (II) except in formula (I) when R_4 is H;~~

20 wherein R' independently of R₁, R₃ or R₄ can be any one of the groups defined above in
21 relation to R₁;
22 R₃ is in α -position and is a hydroxyl group, an acyl-group or an alkoxy group R''O,
23 where R'' independently of R₁, R₃, or R₄ can be any of the groups defined above in
24 relation to R₁;
25 R₄ is in β -position and is hydrogen, ~~an alkyl group~~, an acyl group, or an alkoxy group
26 of the formula R'''O, wherein R''' can be any group mentioned for R₁, independent
27 of R₁, R₂, or R₃, for use as a medicament.

1 2. (Withdrawn)

1 3. (Withdrawn)

1 4. (Withdrawn)

1 5. (Previously Amended) A steroid derivative according to claim 1, wherein
2 said steroid is selected from the group consisting of 5-androstene-3 β ,7 β ,17 α -triol, 5-
3 androstene-3 β ,17 α -diol-7-one, androstane-3 β ,7 β ,17 α -triol and androstane-3 β ,17 α -diol-
4 7-one, or an ester or ether thereof.

1 6. (Withdrawn)

1 7-27. (Cancelled)

1 28. (Withdrawn)

1 29. (New) A steroid derivative according to claim 1, wherein R₁, R', and/or
2 R'' form one or more ether(s) and/or ester(s) with the steroid.



1 30. (New) A steroid derivative according to claim 1 or 29, wherein R₄ is an
2 acyl group, in which hydrogen, or an alkoxy or alkyl group, is attached to the keto
3 group.

1 31. (New) A steroid derivative according to claim 30, wherein R_4 is acetyl
2 (CH_3CO), wherein a keto group is attached to a methyl.

1 32. (New) A steroid derivative of 5-androstene-, 5-pregnenolone or the
2 corresponding saturated derivative capable of interrupting disturbances in Wnt-
3 signaling, cell cycle arrest in G1-phase and/or providing an angiostatic effect.

1 33. (New) A Method for treating disturbances in Wnt-signaling, cell cycle arrest
2 in G1-phase and/or providing an angiostatic effect by administering a therapeutically
3 effective amount of a steroid derivative of 5-androstene-, 5-pregnenolone or
4 corresponding saturated derivatives (androstane- or pregnane-) which steroid derivative
5 is capable of interrupting disturbances in Wnt-signaling, cell-cycle arrest in G1-phase,
6 and/or providing an angiostatic effect thereby treating or preventing a pathologic
7 condition disturbed in this way.

1 34. (New) Method according to claim 33, wherein said steroid derivative is
2 described by formula (I) or (II), the only difference between said formulas being the
3 bond between carbons 5 and 6, as shown below:



6 wherein
7 R_1-O is in the -position and is a hydrogen atom; a protecting group, such as CH_3 ,
8 CH_2OMe , or CH_2O -alkyl;
9 R_2 is $R'O$ in or -position of carbon number 7 or where R_2 is $O=$ or $S=$, where R'
10 independently of R_1 , R_3 , R_4 can be any group mentioned in the definition of R_1 except
11 for hydrogen in formula (I) when R_4 is H, but where R_2 can be hydrogen in formula
12 (II) in all instances;
13 R_3 is in -position and is an hydroxyl-group, an acyl-group or $R''O$, where R''
14 independently can be any group as defined in the above given definition of R_1 , and
15 R_4 is in -position and is hydrogen, an acyl group, or an alkoxy group of the formula
16 $R'''O$, wherein R''' can be any group mentioned under R_1 , independent of R_1 , R_2 or
17 R_3 .

1 35. (New) Method according to claim 34, wherein R_1 , R' and/or R'' form one
2 or more ether(s) and/or ester(s) with the steroid.

1 36. (New) Method according to any one of claims 33-35, wherein one or more
2 pregnane- and/or androstane-derivative corresponding to the steroid is used in the
3 manufacture of the medicament.

1 37. (New) Method according to any one of claims 33-35, wherein said
2 interruption of Wnt-signaling is provided by down-regulating cyclin D1 and/or -
3 catenin.

1 38. (New) Method according to any one of claims 33-35, wherein the steroid
2 derivative is administered in a therapeutically effective amount to prevent and/or
3 counteract overexpression of factors present in the Wnt-signaling pathway in tumour
4 systems with a phenotypic or genotypic deviance in this respect, such as mammary
5 carcinomas, lung cancers, head and neck cancers of squamous cell origin, melanomas,
6 and oesophageal cancers and others.

1 39. (New) Method according to claim 38, wherein said steroid is selected from
2 the group consisting of 17-hydroxy-pregnenolone (17 -OH), 5-androstene-3 ,7 , 17 -
3 triol, 5-androstane-3 , 17 -diol.

1 40. (New) A method for treating non-tumour conditions in the form of
2 neovascularisation or excessive growth of fibroblasts, in the form of hypertrophic scars,
3 keloids, corneal neovascularisation, diabetic retinopathy or exsudative forms of macular
4 degeneration by administering a therapeutically effective amount of a steroid derivative
5 selected from the group consisting of 17-hydroxy-pregnenolone (17 -OH), 5-
6 androstene-3 , 17 -diol, 5-androstene-3 ,7 , 17 -triol, 5-androstane-3 , 17 -diol.

1 41. (New) A method of producing a medicament for the treatment and/or
2 prevention of a benign and/or malignant tumour, comprising the steps of
3 (a) contacting 5-androstene-3 , 17 -diol, a sulfate donor, a sulphotransferase
4 and PAPS to provide 5-androstene-3beta-ol-17alpha-sulfate (17a-AEDS) or
5 producing 17a-AEDS synthetically
6 (b) combining the 17a-AEDS so produced with a suitable carrier; whereby a
7 medicament which is capable of acting as a ligand to peroxisome proliferators
8 activated receptor- (PPAR) is produced or use 17a-AEDS so produced as a
9 water-soluble prodrug easily converted into steroid by sulfatase, which is
10 present in most cells.

1 42. (New) A method according to claim 41, wherein the enzyme is DHEA-
2 sulfotransferase or a phenolsulphotransferase.